

Collective excitation frequencies of a Bose – Einstein condensate with electromagnetically induced $1/r$ attraction

I.E.Mazets,

Ioffe Physico-Technical Institute, 194021 St.Petersburg, Russia,
e-mail: mazets@astro.ioffe.rssi.ru

Abstract

We investigate collective excitations of an atomic Bose-Einstein condensate in a recently discovered regime [D. O'Dell et al., Phys. Rev. Lett. **84**, 5687 (2000)] of a balance between electromagnetically induced $1/r$ attraction and a short-range interatomic repulsion. The corresponding frequencies for monopole, dipole, and quadrupole modes are calculated numerically in the zero temperature case.

PACS numbers: 03.75.Fi, 34.20.Cf, 34.80.Qb.

Theoretical as well as experimental studies of dilute atomic Bose – Einstein condensates (BECs) have made an impressive progress since 1995, when quantum degeneration in alkali metal vapours was reached at the first time [1]. These systems are interesting especially due to a presence of an interaction between atoms weak enough to provide a development of quite precise theories describing the properties of these non-ideal degenerate Bose-gases. The interatomic potentials act on a short range, moreover, at ultralow temperatures they are effectively described by contact pseudo-potentials containing a single parameter, the s -wave scattering length a [2]. Here we consider $a > 0$, i.e. the short-range potential provides mutual repulsion of atoms.

Recently, a new kind of atomic BEC has been proposed in Ref.[3]. Namely, it is shown that the particular configuration of intense off-resonant laser beams gives rise to an effective $1/r$ interatomic attraction between atoms located well within the laser wavelength (since inelastic collisions destroy very rapidly BECs of number densities more than 10^{14} cm^{-3} , a long-wavelength infrared laser has to be considered). The directions and polarizations of different incident laser beams can be chosen in such a way that the static (near-zone) $1/r^3$ component of dipole-dipole interaction energy is effectively "averaged

out”, and only $1/r$ attraction remains. Finally, the two-particle interaction energy including both usual short-range interactions and such an ”electromagnetically induced gravitation” reads as

$$U_{12}(\mathbf{r}) = \frac{4\pi\hbar^2 a}{M} \delta(\mathbf{r}) - \frac{u}{r}, \quad (1)$$

where M is the atomic mass and the long-range interaction constant u is expressed in SI units as [3]

$$u = \frac{11Iq^2\alpha^2}{4\pi\epsilon_0^2 c}.$$

The laser beam intensity and wavenumber are denoted by I and q , respectively; α is the atomic polarizability at the frequency q/c (practically, because the external radiation belongs to the IR range, α is equal to the static polarizability). A useful parameter $a_* = \hbar^2/(Mu)$ can be called the Bohr radius for the gravitation-like potential. Practically, the ”gravitation” is weak enough, so always $a_* \gg a$.

In Ref.[3] it is shown that a self-bound solution of the modified Gross – Pitaevskii equation taking into account the gravitation-like interatomic potential exists. A cloud of atoms located in a radiation field of necessary geometry giving rise to $1/r$ attraction is stable without any confining external potential of magnetic or optical trap. The situation is similar to that of astrophysics where we have stars — objects those are stable due to a balance between gravitation and a Boltzmannian gas pressure. In our case the pressure arises from the two sources: interatomic repulsion due to the short-range potential and the ”quantum pressure” corresponding to the kinetic energy term in the Schrödinger equation.

Let us consider a zero-temperature case. Expressing the BEC order parameter through condensate density n and phase ϕ as usual [2], $\Psi = n \exp(i\phi)$, we rewrite the complex Gross – Pitaevskii equation as a set of two real equations:

$$\frac{\partial n}{\partial t} + \nabla(n\mathbf{v}) = 0, \quad (2)$$

$$M \frac{\partial \mathbf{v}}{\partial t} + \nabla \left(\frac{1}{2} M v^2 - \frac{\hbar^2}{2m\sqrt{n}} \nabla^2 \sqrt{n} + \frac{4\pi\hbar^2 a}{m} n + U_g \right) = 0,$$

where the local ”gravitation potential” is

$$U_g(\mathbf{r}) = -u \int d^3\mathbf{r}' \frac{n(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|}.$$

The velocity field is introduced by the relation $\mathbf{v} = \hbar \nabla \pi / M$.

If $N \gg \sqrt{a_*/a}$ (N is the total number of atoms in the BEC) then the set of Eqs.(2) admits an analytical steady state solution [3]. Indeed, in such a limiting

case the kinetic energy term is small and can be omitted; the "gravitation" is balanced by the s -wave scattering. The density of such a self-bound degenerate bosonic cloud is equal to

$$n_0(\mathbf{r}) = \frac{N}{4R_0^2} \frac{\sin(\pi r/R_0)}{r} \theta(R_0 - r) = 0, \quad (3)$$

where θ is the Heavyside function, $R_0 = \pi\sqrt{a_*a}$ is the radius of the cloud boundary (in Ref.[3] there is an arhythmic error in the numerical coefficient in the expression for R_0).

In the present paper, we calculate the eigenfrequencies of oscillations of such a self-bound cloud. We introduce small perturbations n_1 and $\mathbf{v}_1$ of the density and velocity, respectively: $n = n_0 + n_1$, $\mathbf{v} = \mathbf{v}_1$. procedure of linearizing of Eqs.(2) is standard [4]. We choose the time dependence of perturbation in the form of $\exp(-i\omega t)$. We calculate the eigenfrequencies ω within the Thomas – Fermi approximation, i.e., we neglect the kinetic energy term in the equation for the velocity, After elimination of the variable $\mathbf{v}_1$, we get finally

$$\tilde{\omega}^2 n_1(\tilde{\mathbf{r}}) + \tilde{\nabla} \left[\frac{\sin(\tilde{r})}{\tilde{r}} \tilde{\nabla} \left(n_1(\tilde{\mathbf{r}}) - \frac{1}{4\pi} \int d^3\tilde{\mathbf{r}}' \frac{n_1(\mathbf{r}')}{|\tilde{\mathbf{r}} - \tilde{\mathbf{r}}'|} \right) \right] = 0. \quad (4)$$

Eq.(4) is written using dimensionless quantities

$$\tilde{\mathbf{r}} = \frac{\pi}{R_0} \mathbf{r}, \quad \tilde{\omega} = \omega \sqrt{\frac{R_0^5 M^2}{\pi^4 N \hbar^2 a}}. \quad (5)$$

It is convenient to introduce the "gravitation potential perturbation" variable

$$\xi(\tilde{\mathbf{r}}) = \frac{1}{4\pi} \int d^3\tilde{\mathbf{r}}' \frac{n_1(\mathbf{r}')}{|\tilde{\mathbf{r}} - \tilde{\mathbf{r}}'|}, \quad (6)$$

so that $n_1 = -\tilde{\nabla}^2 \xi$. Then Eq.(4) can be identically transformed to

$$\tilde{\omega}^2 \tilde{\nabla}^2 \xi + \tilde{\nabla} \left[\frac{\sin(\tilde{r})}{r} (\tilde{\nabla}^2 \xi + \xi) \right] = 0. \quad (7)$$

To solve this equation, firstly, we note that the orbital momentum l and its projection m to an arbitrary axis are conserved. So we can set

$$\xi = \xi_l(\tilde{r}) Y_{lm}(\vartheta, \varphi),$$

where Y_{lm} is the spherical harmonic. The only remaining thing now is to set the boundary condition for Eq.(7). They can be derived from the tailoring of the inner and outer solutions at $\tilde{r} = \pi$ (i.e., $r = R_0$). First of all, a perturbation does not change the total number N of atoms. So, for monopole modes, the radial derivative of the potential perturbation vanishes at $\tilde{r} > \pi$. For modes

with $l = 1, 2, 3, \dots$ at the same distances ξ_l decreases as $r^{-(l+1)}$. Hence, the boundary conditions are

$$\frac{d\xi_0}{d\tilde{r}}|_{\tilde{r}=\pi} = 0 \quad (8)$$

for the monopole mode and

$$\frac{1}{\xi_l} \frac{d\xi_l}{d\tilde{r}}|_{\tilde{r}=\pi} = -\frac{l+1}{\pi} \quad (9)$$

for the other modes ($l = 1, 2, 3, \dots$).

We solved Eq.(7) numerically by expansion in series:

$$\xi_l(\tilde{r}) = \sum_k C_{lk} j_l(\gamma_{lk} \tilde{r} / \pi).$$

Here j_l is the spherical Bessel function [5]. To satisfy the boundary conditions, one must choose the coefficient γ_{lk} equal to the k -th root of the equation

$$j_{l'}(\gamma) = 0, \quad (10)$$

where

$$\begin{aligned} l' &= 1, & l &= 0, \\ l' &= l - 1, & l &= 1, 2, 3, \dots \end{aligned}$$

For $l = 1$, the lowest frequency is exactly zero. Indeed, the lowest dipole mode corresponds to a translation of a self-bound BEC as a whole in a space free of any external trap potential.

We calculated numerically the four lowest frequencies for $l = 0, 1$, and 2 . The estimated numerical uncertainty is of order of one per cent. Thus, $\tilde{\omega}$ is equal to

$$\begin{aligned} &0.617, 1.27, 1.85, 2.39, \dots \text{ for } l = 0, \\ &0, 0.898, 1.51, 2.06, \dots \text{ for } l = 1, \\ &0.391, 1.10, 1.71, 2.28, \dots \text{ for } l = 2. \end{aligned}$$

Finally, we note that calculation of quantum depletion in a BEC self-bound by "electromagnetically induced gravitation" could be interesting.

This work is supported by the Russian Foundation for Basic Research (project 99-02-17076) and the State Program "Universities of Russia" (project 015.01.01.04).

References

- [1] M.H. Anderson et al., Science **269**, 198 (1995); C.C. Bradley et al., Phys. Rev. Lett. **75**, 1687 (1995); K.B. Davis et al., Phys. Rev. Lett., **75**, 3969 (1995).

- [2] For a recent review of the atomic BEC theory see, e.g., F. Dalfovo, S. Giorgini, L.P. Pitaevskii, and S. Stringari, Rev. Mod. Phys. **71**, 463 (1999).
- [3] D. O'Dell, S. Giovanazzi, G. Kurizki, and V.M. Akulin, Phys. Rev. Lett. **84**, 5687 (2000).
- [4] S. Stringari, Phys. Rev. Lett. **77**, 2360 (1996).
- [5] M. Abramowitz and I.A. Stegun, eds. "*Handbook of Mathematical Functions*", Nat. Bureau of Standards, 1964, Chapter 10.